



# Firefighters & Melanoma

Funded by the AFG Fire Prevention & Safety Grants





**2022**

**DetecTogether**

**Authors:**

**Walker S. Carlos Poston, Ph.D., M.P.H., FACE**

**Sara A. Jahnke, Ph.D., FACE**

**Maria DH Koeppl, Ph.D.**

**Christopher K. Haddock, Ph.D., PStat<sup>®</sup>**

**Center for Fire, Rescue & EMS**

**Health Research**

**NDRI-USA, Inc.**

**New York, NY**

The project team would like to acknowledge and thank Captain Robert Webb (Ret., Fort Worth Fire Department) who has been committed to supporting firefighters in understanding their cancers and sharing empirical evidence with them in their time of need. His assistance and review were appreciated by the team and his dedication to supporting his brothers and sisters is an inspiration.

## GENERAL EPIDEMIOLOGY: MELANOMA

Skin cancer is the most common form of cancer, although melanoma accounts for only 1% of skin cancer cases, despite being responsible for the majority of skin cancer deaths<sup>1</sup>. In 2021, the American Cancer Society (ACS) estimated 106,110 new cases of melanoma will be diagnosed, and 7,180 individuals will die from it<sup>1</sup>. The lifetime risk for developing melanoma is 1 in 38 for Caucasians, 1 in 1,000 for African Americans, and 1 in 167 for Hispanics<sup>1</sup>. Firefighters have an elevated risk for developing melanoma, and as with many cancers, the stage at diagnosis impacts survival rates. Survival rates for melanoma vary depending on the stage at diagnosis, although the combined 5 year relative survival rate is 93%<sup>2</sup>. Diagnosed at Stage 1, the survival rate is 99%, and diagnosed at Stage 4, the survival rate is 27%.

## INTERNATIONAL AGENCY FOR RESEARCH ON CANCER (IARC)

In June 2022, IARC convened an international meeting of scientists to re-evaluate firefighting as an exposure related to cancer. They determined the literature supports reclassifying **firefighting to a Group 1 carcinogen (carcinogenic to humans) based on “sufficient” evidence**<sup>3</sup>. This is the **highest** classification of exposure only assigned when there is scientific certainty.

Their statement indicated:

***There was also “strong” mechanistic evidence that occupational exposure as a firefighter shows the following key characteristics of carcinogens in exposed humans: “is genotoxic”, “induces epigenetic alterations”, “induces oxidative stress”, “induces chronic inflammation”, and “modulates receptor-mediated effects”.***

Specific to melanoma, IARC **noted “limited” evidence in humans for melanoma as related to firefighting**. While typical use of the word “limited” implies a lack of evidence or support, IARC’s classification with the word limited “means that **a positive association has been observed** between exposure to the agent and cancer but that other explanations for the observations (technically termed “chance”, “bias”, or “confounding”) could not be ruled out with reasonable confidence.” ***It should be noted that IARC criteria and classifications are focused on scientific levels of certainty which are more stringent than those focused on the “weight of the evidence”<sup>4</sup> which is often used in cases of workers compensation.***

## GENERAL RISK FACTORS FOR MELANOMA

There are a number of risk factors associated with developing melanoma:

- **Gender:** Before the age of 50, the risk is higher for women, however, after the age of 50, the risk is higher for men<sup>5</sup>.
- **Age:** While melanoma is one of the more common cancers among younger people, the average age of diagnosis remains approximately 65 years<sup>1</sup>.
- **Race:** Melanoma is more than 20 times more common in Caucasians than in African Americans<sup>1</sup>. This may be related to the risk of melanoma being higher for individuals with red or blonde hair, blue or green eyes, or fair skin<sup>5</sup>.
- **Personal health history:** Having a weakened immune system increases the risk of melanoma, as does having a history of melanoma or other skin cancers<sup>5</sup>, including a family history. Having a first-degree relative (parents, siblings, or children) who has had melanoma increases the likelihood of developing it.
- **Ultraviolet light exposure:** Exposure to UV light is a major risk of melanoma. Examples of exposure include tanning beds, sun lamps, or direct sun exposure<sup>5</sup>.

## RISK FACTORS RELEVANT TO FIREFIGHTERS

Firefighters are exposed to a broad range of chemicals, both in the firehouse and during emergency response. Recent research conducted with live burns has begun to identify and quantify the presence of carcinogens that are typically present on the fire ground. Most alarming are findings that, even when the air appears “clear” there are often ultra-fine respirable particles and gaseous chemicals of several known carcinogens present. Unfortunately, this time period when there is no visible smoke is typically when firefighters remove their personal protective equipment and self-contained breathing apparatus. Particularly noted in the research is the presence of carcinogens such as polybrominated diphenyl ethers (PBDEs), perfluorooctanoic and perfluorooctanesulfonic acids (PFOA and PFOS), phthalates, dioxins, benzene, polychlorinated biphenyls (PCBs), polycyclic aromatic hydrocarbons (PAHs), vinyl chloride, and heavy metals<sup>6-14</sup>. Firefighters face several routes of exposure including inhalation, dermal absorption, secondary exposure through contaminated dust from particulates post incident, and potentially the semi-volatile off-gassing of gear. PCBs specifically have been identified by the International Agency for Research on Cancer (IARC) as having *sufficient evidence* of causing malignant melanoma in humans<sup>15</sup>. While still being evaluated by the IARC process, many of the other known carcinogens found on the fire ground also have been implicated in the development of melanoma<sup>16,17</sup>.

**Polychlorinated biphenyls (PCBs).** PCBs are man-made organic chemicals commonly used as coolants, lubricants in transformers, capacitors, and other electrical equipment. While the chemicals have been banned since the late 1970s due to evidence that they are a probable human carcinogen, they remain in products manufactured prior to the ban and have been found in the fire environment as a product of combustion and are identified as a major contributor to melanoma in the general population<sup>18</sup> and among firefighters<sup>19</sup>.

Other known fireground exposures with a growing literature relating them to malignant melanoma include:

**Polybrominated Diphenyl Ethers (PBDEs).** PBDEs are a complex grouping of chemicals present in polyurethane foam in furniture, electronics, plastics, and flame retardants. Clearly, these common household products are present in many house fires and result exposures on the fireground. These chemicals exert effects on hormonal systems and the thyroid systems, linking them to the development of cancer<sup>20,21</sup> in general and malignant melanoma in particular<sup>15</sup>.

**Benzene.** Benzene is present as a product of combustion from several standard household materials (e.g. PVC pipe, PVC siding, Christmas trees)<sup>6</sup>, from exposure to diesel exhaust, and has been found to off-gas from firefighters’ PPE<sup>10</sup> and is widely recognized as a fireground risk. Benzene is not only present on the fire ground as a product of combustion, but also at high rates in many fire stations as trucks and ambulances are housed in the bay areas. While efforts are being made to increase the use of exhaust mitigation devices in the firehouse, their introduction and use is relatively new to the fire service. Exposure to benzene has been found to increase malignant melanoma risk<sup>16</sup>.

**Shift Work.** In 2019, the International Agency for Research on Cancer classified alternative shift work (including evening, night, rotating, and other unspecified schedules) as a probable human

carcinogen<sup>22</sup>. The relationship between shift work and cancer development occurs through several mechanisms, including circadian rhythm disruptions, impacted melatonin secretion and production, and affecting lifestyle choices<sup>23</sup>. Given the nature of the job and emergency calls, it is not surprising that firefighters – who are faced with a career of 24–48-hour shifts and emergency calls during the night – struggle with the negative health implications of shift work. Long-term shift work, especially night shifts, has been linked to an increased risk of cancer<sup>24–27</sup>, including melanoma<sup>28</sup>.

## RISK OF MELANOMA AMONG FIREFIGHTERS

A number of methodologically sound studies have studied the relationship between melanoma and firefighting and have found increased risks.

For instance, Tsai et al.<sup>29</sup> conducted a case-control study of cancer risk among firefighters in California from 1988-2007 using the California Cancer Registry (CCR). The study included 3,996 male firefighters with cancer and 48,725 non-firefighter controls. **The authors found that firefighters were at a 56% greater risk for melanoma cancer** (OR = 1.56, 95% CI = 1.44-2.13) even after adjusting for age of diagnosis, race, and year of diagnosis.

Lee et al.<sup>30</sup> retrospectively examined over 100,000 career Florida firefighters over a 34-year period, identifying 3,760 male and 168 female primary cancer incidences using the Florida State Fire Marshall's Office and Florida Cancer Data System. After adjusting for age and year of cancer diagnosis, the authors found **male firefighters had 56% higher risk of developing melanoma** (aOR = 1.56, 95% CI = 1.39 – 1.76). While female firefighters had an even higher elevated risk, the relationship was not significant likely due to small sample size (aOR = 1.68, 95% CI = 0.97 – 2.90).

Additional international work has found similar patterns. Pukkala and colleagues<sup>31</sup> examined data from five Nordic countries and found **firefighters had a 25% increased risk** of developing skin cancer compared to the general population (SIR =1.25, 95% CI =1.03 – 1.51). Canadian<sup>32</sup> research found **firefighters were at a 67% greater risk** of developing melanoma (HR=1.67, 95% CI=1.17-2.37), while a cohort study from Australia<sup>33</sup> found **firefighters were 25% more likely to develop melanoma** (SIR=1.25, 95% CI=1.28-1.62). Danish firefighters<sup>34</sup> were found to be **24% more likely** than the general population to develop melanoma (SIR=1.24, 95% CI=0.98-1.57). In a study based on a Canadian cohort of more than a million men, **firefighters were 67% more likely to develop melanoma** (HR=1.67, 95% CI=1.17-2.37) than non-firefighting peers<sup>32</sup>. A comparison of firefighters in Scotland **found a rate 121% higher for melanoma compared to non-firefighters** (13.5 vs. 7.1, 95% CI=3.0-8.8, p<.001)<sup>35</sup>.

**Meta-Analytic Findings.** While individual studies provide a strong foundation for the likely relationship between melanoma and firefighting, even more convincing are the meta-analytic findings. It is difficult to make decisions about whether firefighting increases the risk of any type of cancer based on a single report because results often vary from study to study and studies vary with respect to a number of important design factors including sample size which is a primary determinant of the power to detect group differences. Meta-analysis is a widely-accepted procedure for statistically combining data from multiple studies. *By combining all relevant studies, a meta-analysis provides the maximum possible sample size (at the time the review is conducted) and thus provides the smallest possible confidence interval around an estimate.* This is why meta-analysis is a particular important tool for identifying risks for rare

diseases. Meta-analytic findings for the relationship between melanoma and firefighting provide strong evidence for the relationship

The meta-analysis of LeMasters and colleagues<sup>36</sup> published in 2006 found **more than a doubling of mortality for firefighters from malignant melanoma** (PMR=2.25, 95% CI=1.03-4.27) which led the authors to classify malignant melanoma as probably related to firefighting (*the highest category*).

More recently, a meta-analysis conducted by Jalilian and colleagues<sup>37</sup> found **a 21% elevated risk of developing malignant melanoma** in firefighters (SIREs=1.21, 95% CI=1.02-1.45).

## References

1. Melanoma Skin Cancer Statistics. Accessed November 28, 2021. <https://www.cancer.org/cancer/melanoma-skin-cancer/about/key-statistics.html>
2. Melanoma Survival Rates | Melanoma Survival Statistics. Accessed November 28, 2021. <https://www.cancer.org/cancer/melanoma-skin-cancer/detection-diagnosis-staging/survival-rates-for-melanoma-skin-cancer-by-stage.html>
3. Demers PA, DeMarini DM, Fent KW, et al. Carcinogenicity of occupational exposure as a firefighter. *Lancet Oncol.* 2022;23(8):985-986. doi:10.1016/S1470-2045(22)00390-4
4. Guidotti T. Cancer. In: *Health Risks and Fair Compensation in the Fire Service*. Risk, Systems and Decisions. Springer; 2016.
5. Melanoma Skin Cancer Risk Factors | Melanoma Risk Factors. Accessed November 28, 2021. <https://www.cancer.org/cancer/melanoma-skin-cancer/causes-risks-prevention/risk-factors.html>
6. Fabian T, Borgerson J, Kerber S, et al. *Firefighter Exposure to Smoke Particulates*. Underwriters Laboratories Inc; 2010:6.
7. Baxter CS, Hoffman JD, Knipp MJ, Reponen T, Haynes EN. Exposure of Firefighters to Particulates and Polycyclic Aromatic Hydrocarbons. *J Occup Environ Hyg.* 2014;11(7):D85-D91. doi:10.1080/15459624.2014.890286
8. Fent KW, Eisenberg J, Snawder J, et al. Systemic Exposure to PAHs and Benzene in Firefighters Suppressing Controlled Structure Fires. *Ann Occup Hyg.* 2014;58(7):830-845. doi:10.1093/annhyg/meu036
9. Fent KW, Alexander B, Roberts J, et al. Contamination of firefighter personal protective equipment and skin and the effectiveness of decontamination procedures. *J Occup Environ Hyg.* 2017;14(10):801-814. doi:10.1080/15459624.2017.1334904
10. Fent KW, Evans DE, Booher D, et al. Volatile Organic Compounds Off-gassing from Firefighters' Personal Protective Equipment Ensembles after Use. *J Occup Environ Hyg.* 2015;12(6):404-414. doi:10.1080/15459624.2015.1025135
11. Kirk KM, Logan MB. Firefighting instructors' exposures to polycyclic aromatic hydrocarbons during live fire training scenarios. *J Occup Environ Hyg.* 2015;12(4):227-234. doi:10.1080/15459624.2014.955184
12. Kirk KM, Logan MB. Structural Fire Fighting Ensembles: Accumulation and Off-gassing of Combustion Products. *J Occup Environ Hyg.* 2015;12(6):376-383. doi:10.1080/15459624.2015.1006638
13. Robinson MS, Anthony TR, Littau SR, et al. Occupational PAH Exposures during Prescribed Pile Burns. *Ann Occup Hyg.* 2008;52(6):497-508. doi:10.1093/annhyg/men027
14. Jahnke SA, Jitnarin N, Kaipust CK, Hollerbach BH, Naylor BM, Crisp, C. *Fireground Exposure of Firefighters: A Literature Review*. Fire Protection Research Foundation; 2021.

15. International Agency for Research on Cancer (IARC). *List of Classifications by Cancer Sites with Sufficient or Limited Evidence in Humans, IARC Monographs Volume 1-130.*; 2021. Accessed March 10, 2022. [https://monographs.iarc.who.int/wp-content/uploads/2019/07/Classifications\\_by\\_cancer\\_site.pdf](https://monographs.iarc.who.int/wp-content/uploads/2019/07/Classifications_by_cancer_site.pdf)
16. Mehlman MA. Causal relationship from exposure to chemicals in oil refining and chemical industries and malignant melanoma. *Ann N Y Acad Sci.* 2006;1076:822-828. doi:10.1196/annals.1371.005
17. Falzone L, Marconi A, Loreto C, Franco S, Spandidos DA, Libra M. Occupational exposure to carcinogens: Benzene, pesticides and fibers. *Mol Med Rep.* 2016;14(5):4467-4474. doi:10.3892/mmr.2016.5791
18. Zani C, Ceretti E, Covolo L, Donato F. Do polychlorinated biphenyls cause cancer? A systematic review and meta-analysis of epidemiological studies on risk of cutaneous melanoma and non-Hodgkin lymphoma. *Chemosphere.* 2017;183:97-106. doi:10.1016/j.chemosphere.2017.05.053
19. Brinchmann BC, Bugge MD, Nordby KC, Alfonso JH. Firefighting and melanoma, epidemiological and toxicological associations: a case report. *Occup Med Oxf Engl.* 2022;72(2):142-144. doi:10.1093/occmed/kqab183
20. Zhang Y, Guo GL, Han X, et al. Do Polybrominated Diphenyl Ethers (PBDEs) Increase the Risk of Thyroid Cancer? *Biosci Hypotheses.* 2008;1(4):195-199. doi:10.1016/j.bihy.2008.06.003
21. Steil GJ, Buzzo JLA, de Oliveira Ribeiro CA, Filipak Neto F. Polybrominated diphenyl ethers BDE-47 and BDE-99 modulate murine melanoma cell phenotype in vitro. *Environ Sci Pollut Res Int.* Published online September 17, 2021. doi:10.1007/s11356-021-16455-0
22. IARC. *Night Shift Work.* Vol 124. World Health Organization; 2020. Accessed November 29, 2022. <https://publications.iarc.fr/Book-And-Report-Series/larc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Night-Shift-Work-2020>
23. Wang XS, Armstrong MEG, Cairns BJ, Key TJ, Travis RC. Shift work and chronic disease: the epidemiological evidence. *Occup Med Oxf Engl.* 2011;61(2):78-89. doi:10.1093/occmed/kqr001
24. Papantoniou K, Devore EE, Massa J, et al. Rotating night shift work and colorectal cancer risk in the nurses' health studies. *Int J Cancer.* 2018;143(11):2709-2717. doi:10.1002/ijc.31655
25. Schernhammer ES, Laden F, Speizer FE, et al. Night-Shift Work and Risk of Colorectal Cancer in the Nurses' Health Study. *JNCI J Natl Cancer Inst.* 2003;95(11):825-828. doi:10.1093/jnci/95.11.825
26. Yuan X, Zhu C, Wang M, Mo F, Du W, Ma X. Retraction: Night Shift Work Increases the Risks of Multiple Primary Cancers in Women: A Systematic Review and Meta-analysis of 61 Articles. *Cancer Epidemiol Biomarkers Prev.* 2019;28(2):423-423. doi:10.1158/1055-9965.EPI-18-1085



27. Dun A, Zhao X, Jin X, et al. Association Between Night-Shift Work and Cancer Risk: Updated Systematic Review and Meta-Analysis. *Front Oncol.* 2020;10:1006. doi:10.3389/fonc.2020.01006
28. Yousef E, Mitwally N, Noufal N, Tahir MR. Shift work and risk of skin cancer: A systematic review and meta-analysis. *Sci Rep.* 2020;10(1):2012. doi:10.1038/s41598-020-59035-x
29. Tsai RJ, Luckhaupt SE, Schumacher P, Cress RD, Deapen DM, Calvert GM. Risk of Cancer Among Firefighters in California, 1988–2007. *Am J Ind Med.* 2015;58(7):715-729. doi:10.1002/ajim.22466
30. Lee DJ, Koru-Sengul T, Hernandez MN, et al. Cancer risk among career male and female Florida firefighters: Evidence from the Florida Firefighter Cancer Registry (1981-2014). *Am J Ind Med.* 2020;63(4):285-299. doi:10.1002/ajim.23086
31. Pukkala E, Martinsen JI, Weiderpass E, et al. Cancer incidence among firefighters: 45 years of follow-up in five Nordic countries. *Occup Environ Med.* 2014;71(6):398-404. doi:10.1136/oemed-2013-101803
32. Harris MA, Kirkham TL, MacLeod JS, Tjepkema M, Peters PA, Demers PA. Surveillance of cancer risks for firefighters, police, and armed forces among men in a Canadian census cohort. *Am J Ind Med.* 2018;61(10):815-823. doi:10.1002/ajim.22891
33. Glass DC, Pircher S, Del Monaco A, Hoorn SV, Sim MR. Mortality and cancer incidence in a cohort of male paid Australian firefighters. *Occup Environ Med.* 2016;73(11):761-771. doi:10.1136/oemed-2015-103467
34. Petersen K, Pedersen J, Bonde J, Ebbelhoej N, Hansen J. Long-term follow-up for cancer incidence in a cohort of Danish firefighters. *Occup Environ Med.* Published online 2017. doi:10.1136/oemed-2017-104660
35. Ide CW. Cancer incidence and mortality in serving whole-time Scottish firefighters 1984-2005. *Occup Med Oxf Engl.* 2014;64(6):421-427. doi:10.1093/occmed/kqu080
36. LeMasters GK, Genaidy AM, Succop P, et al. Cancer risk among firefighters: a review and meta-analysis of 32 studies. *J Occup Environ Med Am Coll Occup Environ Med.* 2006;48(11):1189-1202. doi:10.1097/01.jom.0000246229.68697.90
37. Jalilian H, Ziaei M, Weiderpass E, Rueegg CS, Khosravi Y, Kjaerheim K. Cancer incidence and mortality among firefighters. *Int J Cancer.* 2019;145(10):2639-2646. doi:https://doi.org/10.1002/ijc.32199



**DetecTogether**